

Small molecule discovery

NCATS

COLLABORATE. INNOVATE. ACCELERATE.

Small Molecule Discovery in Oncology and Beyond: Challenges and Opportunities

Anton Simeonov, Ph.D.

*Scientific Director, National Center for Advancing Translational
Sciences (NCATS), National Institutes of Health (NIH)*

**TRACO Lecture
October 10, 2023**




NIH

National Center
for Advancing
Translational Sciences


Drug discovery steps

Steps in the Drug Discovery Process




Screening assays

Range of Screening Assays




Robotics


Robotics & Informatics




Microliter Dispensing



Laser Cytometry




Microscopy



HTS assays

149 Cancer Relevant Cell-Based HTS Assays from PubChem



Coussens, N. P., Braisted, J. C., Peryea, T., Sittampalam, S. G., Simeonov, A. and Hall, M. D. **Small Molecule Screens: A Gateway to Cancer Therapeutic Agents with Case Studies of FDA-Approved Drugs** *Pharmacological Reviews*, October 2017, 69 (4) 479-496



Assay expense

Important Considerations for Choosing an Assay

- **Assay expense**
 - Cost per well
 - Disposal cost(s)

Instrumentation

Important Considerations for Choosing an Assay

- Assay expense
 - Cost per well
 - Disposal cost(s)
- **Available instrumentation**
 - Select the best possible assays based on the available instrumentation

Throughput


Important Considerations for Choosing an Assay

- **Assay expense**
 - Cost per well
 - Disposal cost(s)
- **Available instrumentation**
 - Select the best possible assays based on the available instrumentation
- **Assay throughput**
 - Miniaturization reduces the cost per well


Miniaturization

Assay Miniaturization Saves Time and Reagents



96-well
~50,000 cells/well



384-well
~5,000 cells/well



1536-well
~500 cells/well



	96	384	1536
Plates per 100,000 compounds:	1,042	261	66
Assay volume (uL):	50-200	30-50	2-8
Adherent cell seeding density:	~10,000	~2,000	~500

Horman, Shane R. "Complex High-Content Phenotypic Screening." Special Topics in Drug Discovery. InTech, 2016.

Multiplex

Important Considerations for Choosing an Assay

- **Assay expense**
 - Cost per well
 - Disposal cost(s)
- **Available instrumentation**
 - Select the best possible assays based on the available instrumentation
- **Assay throughput**
 - Miniaturization reduces the cost per well
- **Ability to multiplex**
 - Can the response be measured by a single parameter; is multiparametric output possible?
 - Increased data per sample
 - Can guide hit selection by differentiating selectivity among related targets
 - Can distinguish pathway inhibition from cytotoxicity in a cell-based assay

Reagents

Important Considerations for Choosing an Assay

- **Assay expense**
 - Cost per well
 - Disposal cost(s)
- **Available instrumentation**
 - Select the best possible assays based on the available instrumentation
- **Assay throughput**
 - Miniaturization reduces the cost per well
- **Ability to multiplex**
 - Can the response be measured by a single parameter; is multiparametric output possible?
 - Increased data per sample
 - Can guide hit selection by differentiating selectivity among related targets
 - Can distinguish pathway inhibition from cytotoxicity in a cell-based assay
- **Reagents**
 - **Stability for hours is important**
 - **Consistency is critical (ideally obtain a large quantity from a single lot)**
 - **All reagents need to be validated (cell lines, antibodies, enzymatic purity, etc.)**


Interference

Important Considerations for Choosing an Assay

- **Assay expense**
 - Cost per well
 - Disposal cost(s)
- **Available instrumentation**
 - Select the best possible assays based on the available instrumentation
- **Assay throughput**
 - Miniaturization reduces the cost per well
- **Ability to multiplex**
 - Can the response be measured by a single parameter; is multiparametric output possible?
 - Increased data per sample
 - Can guide hit selection by differentiating selectivity among related targets
 - Can distinguish pathway inhibition from cytotoxicity in a cell-based assay
- **Reagents**
 - Stability for hours is important
 - Consistency is critical (ideally obtain a large quantity from a single lot)
 - All reagents need to be validated (cell lines, antibodies, enzymatic purity, etc.)
- **Potential for assay interference**
 - **Fluorescent compounds can interfere with fluorescent readouts**
 - **Colored compounds might interfere with luminescence**

Spectroscopic profiling



Fluorescence Spectroscopic Profiling of Compound Libraries



Simeonov, A., Jadhav, A., Thomas, C.J., Wang, Y., Huang, R., Southall, N.T., Shinn, P., Smith, J., Austin, C.P., Auld, D.S. and Inglese, J., 2008. **Fluorescence spectroscopic profiling of compound libraries.** *Journal of Medicinal Chemistry*, 51(8), 2363-2371.

Tolerance to DMSO

Determination of Assay Tolerance to DMSO/Vehicle is Important



Yasgar A., Jadhav A., Simeonov A., Coussens N.P., **AlphaScreen-Based Assays: Ultra-High-Throughput Screening for Small-Molecule Inhibitors of Challenging Enzymes and Protein-Protein Interactions.** *Methods Mol Biol.* 2016;1439:77-98.

Homogenous format

Important Considerations for Choosing an Assay

- **Homogenous assay format is preferred for screening**
 - Add reagents, mix and measure (no solution removal or wash steps)
 - Automation friendly
 - Reduces variability
 - Decreases hands-on time
 - Improves reproducibility

Time required

Important Considerations for Choosing an Assay

- Homogenous assay format is preferred for screening
 - Add reagents, mix and measure (no solution removal or wash steps)
 - Automation friendly
 - Reduces variability
 - Decreases hands-on time
 - Improves reproducibility
- **Time required for assay**
 - Off-line reagent preparation
 - Is temperature equilibration required
 - Actual assay time
 - Kinetic versus end point read
 - Time required for data analysis and record keeping

Signal stability

Important Considerations for Choosing an Assay

- **Homogenous assay format is preferred for screening**
 - Add reagents, mix and measure (no solution removal or wash steps)
 - Automation friendly
 - Reduces variability
 - Decreases hands-on time
 - Improves reproducibility
- **Time required for assay**
 - Off-line reagent preparation
 - Is temperature equilibration required
 - Actual assay time
 - Kinetic versus end point read
 - Time required for data analysis and record keeping
- **Signal stability**
 - Does the response occur rapidly or within a few minutes or hours?
 - Longer signal stability allows for flexibility in automated systems
 - Longer signal stability minimizes differences among plates within a stack


Sensitivity

Important Considerations for Choosing an Assay

- **Homogenous assay format is preferred for screening**
 - Add reagents, mix and measure (no solution removal or wash steps)
 - Automation friendly
 - Reduces variability
 - Decreases hands-on time
 - Improves reproducibility
- **Time required for assay**
 - Off-line reagent preparation
 - Is temperature equilibration required
 - Actual assay time
 - Kinetic versus end point read
 - Time required for data analysis and record keeping
- **Signal stability**
 - Does the response occur rapidly or within a few minutes or hours?
 - Longer signal stability allows for flexibility in automated systems
 - Longer signal stability minimizes differences among stacks of plates
- **Assay Sensitivity**
 - Choice of readouts is important
 - Colorimetric < fluorescent < luminescent


Evaluating assay

Evaluating Assay Suitability for Screening



$$S/N = \frac{\text{mean signal} - \text{mean background}}{\text{standard deviation of background}}$$

$$S/B = \frac{\text{mean signal}}{\text{mean background}}$$




$$Z = 1 - \frac{3SD \text{ of sample} + 3SD \text{ of control}}{|\text{mean of sample} - \text{mean of control}|}$$

A Simple Statistical Parameter for Use in Evaluation and Validation of High Throughput Screening Assays. Zhang JH, Chung TD, Oldenburg KR. J Biomol Screen. 1999;4(2):67-73.

Suitability

Evaluating Assay Suitability for Screening




$$Z = 1 - \frac{3SD \text{ of sample} + 3SD \text{ of control}}{|\text{mean of sample} - \text{mean of control}|}$$

Z-factor value	Structure of assay	Related to screening
1	SD = 0 (no variation), or the dynamic range $\rightarrow \infty$	An ideal assay
$1 > Z \geq 0.5$	Separation band is large	An excellent assay
$0.5 > Z > 0$	Separation band is small	A double assay
0	No separation band, the sample signal variation and control signal variation bands touch	A "yes/no" type assay
<0	No separation band, the sample signal variation and control signal variation bands overlap	Screening essentially impossible

A Simple Statistical Parameter for Use in Evaluation and Validation of High Throughput Screening Assays. Zhang JH, Chung TD, Oldenburg KR. J Biomol Screen. 1999;4(2):67-73.

Quantitative HTS

Improving the Process of Early Discovery: Quantitative High-Throughput Screening (qHTS)



- Conventional screening done at one concentration
 - Not appropriate for potency testing – “dose makes the poison”
- qHTS tests compounds assayed at **multiple** concentrations (range: 4 logs)
- Enabled by miniaturized assay volumes (2-8 μL per test) and informatics pipeline
- Generates *pharmacological actives* instead of statistical “hits”
 - Dramatically increases reliability
 - Dramatically reduces false positives and false negatives
- *To date, several hundred million datapoints from several hundred screens have been generated and deposited in the public domain.*

PNAS 103:11473

Medicinal chemistry

Medicinal Chemistry, an Integrated Process



Tier 1: Synthesis & validation



- Medicinal chemistry
- Purification
- In vitro ADME

Tier 2: Compound profile expansion

- Met ID/CYP studies
- In vitro toxicology


Tier 3: Advanced Preclinical studies

- Formulation
- Scale-up
- In vivo PK/PD
- Preclinical toxicology





Drug combinations

Translation Challenge: Rapid Discovery of Drug Combinations



Resistance

Application of Drug Combinations to Address Resistance



Dissemination

Dissemination of technology: combination screening to overcome drug resistance in cancer cells

- Applied to ABC subtype of Diffuse Large B-Cell Lymphoma (ABC-DLBCL)
- Ibrutinib is a BTK inhibitor that has activity against ABC DLBCL
- Lead investigators: Craig Thomas (NCATS) and Louis Staudt (NCI)
- Study evaluated 459 drugs *in combination* with Ibrutinib
 - » 6 x 6 concentration-response “matrix blocks”, validation in 10 x 10 concentration-response matrix blocks
- DNA-damaging agents identified as synergizing with Ibrutinib in killing ABC DLBCL cell lines

Dissemination:



» Protocols

» Source code for dispense

High-throughput combinatorial screening identifies drugs that cooperate with ibrutinib to kill activated B-cell–like diffuse large B-cell lymphoma cells

Leahy A, Matthews Grimer^{1,2}, Rajanishi Guha^{1,2}, Paul Shiner^{1,2}, Ryan M. Young^{1,2}, Jonathan M. Kaffer^{1,2}, Dongbo Liu^{1,2}, Tom S. Goldlust^{1,2}, Adam Yang^{1,2}, Crystal McKnight^{1,2}, Matthew B. Rosen^{1,2}, Damien Y. Duvenyi^{1,2}, Bao-Kiang Jiang^{1,2}, Sam Mitchell^{1,2}, Tim Mirzavani^{1,2}, Willemse Huang^{1,2}, Martin J. Walsh^{1,2}, Bryan T. Miller^{1,2}, Parasoma Patel^{1,2}, William Leiber^{1,2}, David J. Maloney^{1,2}, Christopher A. Leclair^{1,2}, Ganisha Rai^{1,2}, Ajit Jadhav^{1,2}, Brian D. Fryer^{1,2}, Christopher P. Austin^{1,2}, Scott E. Martin^{1,2}, Anton Simionov^{1,2}, Matt Farmer^{1,2}, Louis M. Staudt^{1,2}, and Craig J. Thomas^{1,2}



¹Division of Preclinical Innovation, National Institutes of Health-Chemical Genomics Center, National Center for Advancing Translational Sciences, Metabolism Branch, Center for Cancer Research, and ²Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, and ³Basic Science Program, Merck Research, Inc., Chemical Biology Laboratory, President National Laboratory for Cancer Research, Merck, Kenilworth, NJ 07033



Triple drug combination

Example: triple drug combination screening to tackle resistance against artemisinin-based combination therapies in malaria




ACS Pharmacol. Transl. Sci. 2020, <https://dx.doi.org/10.1021/acscptsci.0c00110?ref=pdf>



- ① Drugs A and B are acoustically dispensed in a 10x10-well matrix, 12 replicate blocks per plate. Single drug responses, bottom row (Drug A) and right column (Drug B).

- ② To each replicate block, serial dilutions of Drug C is acoustically dispensed, with the final block serving as a DMSO control

- ③ Plate view of triple combination screening plate with positive control (artesunate, blue) and neutral controls (DMSO, grey) also shown.



- ④ Dispense *P. falciparum* and erythrocytes, incubate 72 hr

- ⑤ Dispense 2 μ L of SYBRGreen1 and lysis solution, incubate overnight. Fluorescence quantified

- ⑥ Parasite proliferation response is normalized to artesunate and DMSO controls. For each concentration Drug C block, response of Drug A + Drug B wells is summed.

- ⑦ Triple drug response is analyzed as a function of Drug C concentration.

VIPOR combination

NIH U.S. National Library of Medicine
ClinicalTrials.gov

Find Studies ▾ About Studies ▾ Submit Studies ▾ Resources ▾ About Site ▾ [PRS Login](#)

[Home](#) > [Search Results](#) > Study Record Detail Save this study

Venetoclax, Ibrutinib, Prednisone, Obinutuzumab, and Revlimid (VIPOR) in Relapsed/Refractory B-cell Lymphoma

⚠ The safety and scientific validity of this study is the responsibility of the study sponsor and investigators. Listing a study does not mean it has been evaluated by the U.S. Federal Government. [Know the risks and potential benefits](#) of clinical studies and talk to your health care provider before participating. Read our [disclosure](#) for details.

ClinicalTrials.gov Identifier: NCT03223610

Recruitment Status: **Recruiting**
First Posted: July 21, 2017
Last Update Posted: July 7, 2022
[See Contacts and Locations](#)

Sponsor:

National Cancer Institute (NCI)

Information provided by (Responsible Party):

National Institutes of Health-Clinical Center (CC) (National Cancer Institute (NCI))

VIPOR Regimen Is Safe, Shows Impressive Activity in Relapsed/Refractory DLBCL

November 14, 2020

Onco News



December 23, 2020 — The 5-drug regimen of venetoclax, ibrutinib, prednisone, obinutuzumab, and lenalidomide showed a tolerable safety profile and encouraging antitumor activity with complete responses in patients with relapsed/refractory diffuse large B-cell lymphoma.



Christopher J. Horne, MD

The 5-drug regimen of venetoclax (Venclexta), ibrutinib (Imbruvica), prednisone, obinutuzumab (Gazyva), and lenalidomide (Revlimid; VIPOR) showed a tolerable safety profile and encouraging antitumor activity with complete responses (CRs) in patients with relapsed/refractory diffuse large B-cell lymphoma (DLBCL), according to phase 1b/2 findings that were presented during the 2020 ASH Annual Meeting and Exposition.

Results showed that of 52 evaluable patients, the best overall response was 71%, with a complete response (CR) rate of 52% and a 19% partial response (PR) rate. Specifically, in relapsed patients (n = 30), the objective response rate (ORR) was 50% with a 70% and 13% CR and PR rate, respectively. The ORR was 55% in refractory patients (n = 22), with a 27% CR rate and a 27% PR rate.

<https://www.onclive.com/view/vipor-regimen-is-safe-shows-impressive-activity-in-relapsed-refractory-dlbcl>

VIPOR Regimen Signals Benefit in Patients With Mantle Cell Lymphoma

February 7, 2022

Onco Today



In an interview with Targeted Oncology, Christopher Horne, MD, discussed the ongoing VIPOR study exploring a 5-drug regimen of venetoclax and CD20 antibodies, and NP-1 and anti-CD20 antibody combination.

Treatment with the VIPOR regimen—consisting of venetoclax (Venclexta), ibrutinib (Imbruvica), prednisone, and lenalidomide (Revlimid), has thus far appeared safe for use in patients with mantle cell lymphoma (MCL) and has demonstrated preliminary activity.

Results from the phase 1 portion of the VIPOR study (NCT03223610) were presented during the 63rd American Society of Hematology (ASH) Annual Meeting & Exposition. Of the 11 patients who were treated, the ORR was 100% and the complete remission (CR) rate was 80%. Nine patients were evaluable for safety and no dose-limiting toxicities were observed. There were few grade 3 and 4 adverse events (AEs), but the hematologic grade 3/4 AEs included neutropenia (13%), anemia (11%), and thrombocytopenia (3%). The non-hematologic grade 3/4 AEs included hypokalemia (33%) along with fatigue, hypomagnesemia, elevated bilirubin, atrial fibrillation, lung infection, and syncope occurring in 11% of patients each.



Christopher Horne, MD

<https://www.targetedonc.com/view/vipor-regimen-signals-benefit-in-patients-with-mantle-cell-lymphoma>



NIH National Center for Advancing Translational Sciences



3D models

Increasing the predictivity of *in vitro* assays: a continuum of 3D models of healthy and diseased tissues




Tissue bioprinting

3D Tissue Bioprinting



Hydrogel polymer is mixed with cells and loaded into syringe.




The printer “3D prints” the cell/gel mixture in a layer by layer approach.




Printed construct



1 day



1 week




2 weeks

The printed construct is incubated to allow the cells to form a tissue, and to enable proper cell differentiation.

3D tissue models

Current portfolio of engineered 3D tissue models



Program Director Marc Ferrer, Ph.D. <https://ncats.nih.gov/bioprinting>

Skin biofabrication

Skin biofabrication

Native Skin



<http://www.siumed.edu/~dking2/intro/IN005b.htm>

3D-Bioprinted Skin

Stratum corneum

Stratum granulosum

Stratum spinosum

Stratum germinativum

Dermis:
Collagen and elastic
fibers



Skin tissue generation

Generation of bioprinted skin tissues


Reconstructed human epidermis (RhE)

1. Coat the 96-well transwell insert membrane with collagen

2. Add keratinocytes

3. Submerge culture for 3 days

4. Air-liquid interface culture for 8 days



2. Bioprint fibroblast bioink to a 3-layer U shape on bottom side of 96-well transwell insert membrane


4. Submerge bioprinted tissue in medium for 7 days

5. Add keratinocytes and submerge culture for 3 days


6. Air-liquid interface culture for 8 days

Full thickness skin tissue (FTS)

1. Suspend fibroblasts in bioprinting gel



3. Add bioprinting gel to cover the U shape




Tissue model

www.oncotarget.com

Oncotarget, 2020, Vol. 11, (No. 27), pp: 2587-2596

Research Paper

A 3D biofabricated cutaneous squamous cell carcinoma tissue model with multi-channel confocal microscopy imaging biomarkers to quantify antitumor effects of chemotherapeutics in tissue



Collaboration between NCATS (Marc Ferrer) and Rockefeller University (Daniel Gareau)


Information



**Where do I go for more
information about assay
development?**

Book

Sharing internal know-how: Assay Guidance Manual (47 chapters/ 1,338 printed pages)



<https://ncats.nih.gov/agm-video>

August 7th Videos

1. Austin, CP: Welcome to the Assay Guidance Manual (AGM) Workshop
2. Coussens, NP: Strategies for Assay Selection & Robust Biochemical Assays
3. Riss, T: Treating Cells as Reagents to Design Reproducible Screening Assays
4. Trask, OJ: Assay Development Considerations for High Content Imaging
5. Auld, DS: Studies in Mechanisms and Methods in Assay Interferences
6. Dahlin, JL: Assay Interference by Chemical Reactivity
7. Chung, TDY: Basic Assay Statistics, Data Analysis & Rules of Thumb
8. Devanarayan, V: Reproducibility & Differentiability of Potency Results
9. Sittampalam, GS: Avoiding Artifacts & Interferences in Assay Operations

March 26-27th Videos

1. Austin, CP: Welcome to the Assay Guidance Manual (AGM) Workshop
2. Coussens, NP: Robust Assays Define Success in Preclinical Research
3. Lal-Nag, M: Target Identification & Validation in Translational Discovery
4. Foley, TL: Development & Validation of Cell-Based and Biochemical Assays
5. Riss, T: Treating Cells as Reagents to Design Reproducible Screening Assays
6. Trask, OJ: Assay Development for HCS & Best Practices for 3D HCS
7. Roth, KD: Mass Spectrometry for Drug Screening and Lead Optimization
8. Dahlin, JL: Bioassay Interference by Aggregation and Chemical Reactivity
9. Patnaik, S: Lead Selection and Optimization by Medicinal Chemistry
10. Xia, M: *In Vitro* Toxicological Testing Using a qHTS Platform
11. Xu, X: *In Vitro* Assessment of ADME Properties of Lead Compounds
12. Kahl, SD: Statistical Design of Experiments for Assay Development
13. Guha, R: Pharos Application to Target Evaluation and Drug Discovery
14. Weidner, JR: Assay Operations: Keeping Assays Robust and Reproducible

Table of Contents

Preface	
Considerations for Early Phase Drug Discovery	1 Chapter
<i>In Vitro</i> Biochemical Assays	10 Chapters
<i>In Vitro</i> Cell Based Assays	19 Chapters
<i>In Vivo</i> Assay Guidelines	2 Chapters
Assay Artifacts and Interferences	4 Chapters
Assay Validation, Operations and Quality Control	5 Chapters
Assay Technologies	2 Chapters
Instrumentation	2 Chapters
Pharmacokinetics and Drug Metabolism	1 Chapter
Glossary of Quantitative Biology Terms	1 Chapter

Website: <https://ncats.nih.gov/expertise/preclinical/agm>

Email us: NCATS_AGM_Editors@mail.nih.gov



Facebook: www.facebook.com/assayguide



LinkedIn: www.linkedin.com/groups/7427344



NIH National Center
for Advancing
Translational Sciences

Assay guidance manual

Assay Guidance Manual Training Workshops

- [Online Training Modules](#)
- [Upcoming Workshops](#)
- [Past Workshops](#)


NCATS offers a variety of [Assay Guidance Manual \(AGM\)](#) training workshops throughout the year designed to share best practices and advice on robust assay design, development and implementation for researchers involved in the drug discovery process.



Online Training Modules

NCATS offers an online AGM training workshop in addition to the in-person AGM workshops held throughout the year. The online training workshop also features experts sharing best practices and expert advice on assay design, development and implementation. [View the video modules.](#)

<https://ncats.nih.gov/expertise/preclinical/agm/training>



Assay Guidance Workshop on 3D Tissue Models for Antiviral Drug Development

🕒 Tue Jun 7, 11:00 AM - Wed Jun 8, 5:15 PM (EDT)

📍 Zoom

[Add to calendar](#)

[Share](#)

THIS EVENT HAS ENDED

Video of the workshop is available at the links below:

[Assay Guidance Workshop on 3D Tissue Models for Antiviral Drug Development \(Day 1\) \(June 7, 2022\)](#)


[Assay Guidance Workshop on 3D Tissue Models for Antiviral Drug Development \(Day 2\) \(June 8, 2022\)](#)

About the Workshop

The National Center for Advancing Translational Sciences (NCATS) Assay Guidance Manual (AGM) program is hosting a two-day workshop that will cover a broad range of critical concepts, including practical approaches and best practices, for developing standardized 3D cellular assays with the hope of helping the community to successfully develop therapeutics for future pandemic threats. This workshop is jointly organized by NCATS, the [National Institute of Allergy and Infectious Diseases \(NIAID\)](#) and the [Bill & Melinda Gates Foundation](#). The overall goal of this workshop is to help scientists establish robust, reproducible, scalable, consistent, advanced 3D tissue models to study pandemic threat viruses.

Biomedical changes


Changes in Biomedical Research Fields in the Last 100 Years



Joining forces

Chemistry, Engineering, Informatics, and Biology Joining Forces to Automate the Design, Generation, and Testing of New Molecules

A Specialized Platform for Innovative Research Exploration (ASPIRE)





Biological assay development

Original Research

Cross-Platform Bayesian Optimization System for Autonomous Biological Assay Development

SLAS Technology
2021, Vol. 26(6) 579-590
© Society for Laboratory
Automation and Screening 2021
DOI: 10.1177/104726902110053792
journals.sagepub.com/home/jla
SAGE

Sam Elder^{1*}, Carleen Klumpp-Thomas^{2*}, Adam Yasgar², Jameson Travers²,
Shayne Frebert², Kelli M. Wilson² , Alexey V. Zakharov², Jayme L. Dahlin² ,
Christoph Kreisbeck¹, Dennis Sheberla¹, Gurusingham S. Sittampalam²,
Alexander G. Godfrey², Anton Simeonov², and Sam Michael²



Graphical Abstract.

NCATS

NCATS

COLLABORATE. INNOVATE. ACCELERATE.

 ncats.nih.gov

 [@ncats_nih_gov](https://twitter.com/ncats_nih_gov)

 [@ncats.nih.gov](https://facebook.com/ncats.nih.gov)

 [NIH-NCATS](https://linkedin.com/company/NIH-NCATS)